


Chateau® Herbicide SW—A Proven Preemergence Residual Herbicide—Now for Citrus

- ▶ **Broad Spectrum:** controls more than 90 annual broadleaf and grass weeds, including spurge, eclipta, Florida pusley, nightshades, marestail, fleabane, mallow, filaree and many small seeded winter annual weeds
- ▶ **Long-Lasting:** stays where you put it; won't leach or co-distillate; no photodegradation
- ▶ **Grove-Ready:** whether newly-planted or established, *Chateau* is effective and remains active until ¼-inch rainfall or irrigation releases its activity



Chateau Delivers Goatweed Control

4 months after application



Means followed by the same letters are not significantly different

Source: Dr. Brent Sellers, University of Florida

How To Use

Rate	<ul style="list-style-type: none"> • 6–12 oz/sprayed acre. For maximum residual control, use the higher rate. Apply with a burndown herbicide if emerged weeds are present. • If weed pressure is heavy, make two sequential applications. Wait at least 30 days between applications. • Maximum 24 oz/A per year
Method	Spray a uniform band directed at the base of the trunk
Spray Volume	10–30 GPA
Tank Mix Partners	Add burndown products to control emerged weeds (Cheetah®, glyphosate, Gramoxone Inteon®, Treevix®, Rely® (glufosinate), etc.). <i>Chateau</i> can also be tank mixed with other preemergence herbicides, such as Prowl® or Alion® to extend weed spectrum following most-restrictive labeling.
REI / PHI	12 hours / 3 days

Chateau Delivers Control of Common Citrus Weeds



Annual Fleabane Control

Effective, long-lasting control of water and nutrient-robbing annual fleabane.

Source: Valent

Horseweed Control

Control horseweed (marestail) before it emerges. If established weeds are present combine with an effective postemergence product, such as Roundup PowerMAX.

Source: Valent

Other Important Information

- ▶ A maximum rate of 6 oz/A per application should be used on any soil that has a sand plus gravel content over 80% if trees are less than 3 years of age. (Two applications of 6 oz/A in a 12-month period can still be made as long as there have been 60 days between applications.)
- ▶ Apply carefully—although *Chateau* is primarily a preemergence herbicide, direct spray contact or drift will damage foliage, green bark and fruit
- ▶ Do not apply to trees established <1 year unless protected from spray with non-porous wrap, grow tubes or waxed containers
- ▶ Do not apply to farm alleys or roads where traffic may result in treated dust settling onto crops or other desirable vegetation
- ▶ If using a spray drift agent with all *Chateau* applications, add ammonium sulfate (AMS) to spray tank before mixing *Chateau* and a spray drift agent. Conduct a jar test before mixing.
- ▶ Clean out spray tank daily